



E573
JACC March 12, 2013
Volume 61, Issue 10

Heart Failure

THE RELATIONSHIP BETWEEN TRICUSPID ANNULAR SYSTOLIC EXCURSION AND PULMONARY ARTERIAL SYSTOLIC PRESSURE IN HEART FAILURE: AN INDICATOR OF CARDIAC LENGTH VERSUS FORCE RELATIONSHIP THAT DISCLOSES DISEASE SEVERITY AND PROGNOSIS

Oral Contributions

West, Room 2006

Saturday, March 09, 2013, 9:15 a.m.-9:30 a.m.

Session Title: Resynchronization Therapy and Right Ventricular Function

Abstract Category: 15. Heart Failure: Clinical

Presentation Number: 906-8

Authors: *Francesco Bandera, Ross Arena, Serenella Castelveccchio, Lorenzo Menicanti, Valentina Labate, Greta Generati, Eleonora Alfonzetti, Stefano Ghio, Pier Temporelli, Marco Guazzi, Heart Failure Unit, Cardiology, University of Milano, IRCCS Policlinico San Donato, San Donato Milanese, Milano, Italy*

Background: Right ventricular (RV) dysfunction carries a remarkable risk of death in heart failure (HF). Echo-derived pulmonary artery systolic pressure (PASP) and tricuspid annular systolic excursion (TAPSE), as an indicator of RV systolic function, are of remarkable clinical value in the follow-up of HF patients. We hypothesized that the relationship between TAPSE (longitudinal RV fiber shortening) and PASP (developed force) would provide an index of RV force-length relationship and that their ratio would better disclose disease severity and prognosis compared to either variable independently.

Methods: A cohort of 293 stable HF patients with both reduced (HFrEF, n=247) and preserved left ventricular ejection fraction (HFpEF, n=46) underwent echo-Doppler and tissue Doppler imaging as well as NT-pro-BNP biomarker assessment and prospectively tracked for adverse events.

Results: TAPSE vs PASP relationship showed a downward shift of the regression line for nonsurvivors that were distributed in the more unfavorable relationship with highest PASP and lower TAPSE. A similar regression line was observed for patients with HFrEF and HFpEF. Given the collinearity between both TAPSE and PASP and the TAPSE/PASP ratio, separate Cox regression and Kaplan-Meier analyses were performed; the first with TAPSE and PASP as individual measures and the second combining them in ratio form. HR for variables retained in the multivariate regression, using the TAPSE/PASP ratio as optimal dichotomous threshold were: TAPSE/PASP \leq 0.36 (HR: 10.4, $p < 0.01$); NT-pro-BNP \leq 1060 pg/ml (HR: 8.9 $p < 0.001$); TAPSE \leq 16 mm (HR: 5.1 $p < 0.01$); NYHA \leq 3 (HR: 4.4, $p < 0.01$); E/E' (HR: 4.1; $p < 0.01$).

Conclusions: This study shows that the TAPSE vs PASP relationship is shifted downward in non-survivors, discloses disease severity and prognosis better than individual variable assessment and is similarly distributed in HFrEF and HFpEF. Results confirm that echo-derived RV systolic function assessment is of basic relevance in HF. The TAPSE vs PASP as a possible index of the length/force relationship is a step forward for a more efficient RV function evaluation and is not affected by the quality of left ventricular dysfunction.